

High-Coverage Whole-Exome Sequencing Identifies Candidate Genes for Suicide in Victims with Major Depressive Disorder

Dóra Tombác^{1,2,†}, Zoltán Maróti^{3,†}, Tibor Kalmár³, Zsolt Csabai¹, Zsolt Balázs¹, Shinichi Takahashi², Miklós Palkovits⁴, Michael Snyder^{2*}, Zsolt Boldogkői^{1*}

Addresses:

¹Department of Medical Biology, Faculty of Medicine, University of Szeged, Somogyi B. u. 4., Szeged, H-6720, Hungary

²Department of Genetics, School of Medicine, Stanford University, 300 Pasteur Dr., Stanford, CA 94305-5120, USA

³Department of Paediatrics, Faculty of Medicine, University of Szeged, Somogyi B. u. 4., Szeged, H-6720, Hungary

⁴Neuromorphological and Neuroendocrine Research Laboratory, Department of Anatomy, Histology and Embryology, Semmelweis University, Budapest, Üllői u. 26., H-1085, Hungary

[†]These authors contributed equally to this work.

*Co-corresponding authors

E-mails:

tombacz.dora@med.u-szeged.hu

maroti.zoltan@med.u-szeged.hu

kalmar.tibor@med.u-szeged.hu

csabai.zsolt@med.u-szeged.hu


balazs.zsolt@med.u-szeged.hu

palkovits.miklos@med.semmelweis-univ.hu

takahashi.shinichi.dz@daichisankyo.co.jp

mepsnyder@stanford.edu

boldogkoi.zsolt@med.u-szeged.hu



S1 Figure. Gene network of candidate genes based on experimental evidence of interaction. The gene network was constructed from all the implicated genes (S4 Table) using the string database accessible at <http://string-db.org/> (version 10.5). All edges represent experimental evidence of interaction. Confidence cut-off was set to 0.333.

